arXiv:0804.0964v1 [math.AC] 7 Apr 2008

Asymptotic stability of certain sets of associated prime ideals of local cohomology modules

Nguyen Tu Cuong[a], Nguyen Van Hoang[b] and Pham Huu Khanh[c]

Institute of Mathematics
18 Hoang Quoc Viet Road, 10307 Hanoi, Vietnam

Abstract[1]. Let $(R, \mathfrak{m})$ be a Noetherian local ring I, J two ideals of R and M a finitely generated $R-$module. It is first shown that for $k \geq -1$ the integer $r_k = \operatorname{depth}_k(I, J^nM/J^{n+1}M)$, it is the length of a maximal $(J^nM/J^{n+1}M)-$sequence in dimension $> k$ in I defined by M. Brodmann and L. T. Nhan [4], becomes for large n independent of n. Then we prove in this paper that the sets $\bigcup_{j\leq r_k} \operatorname{Ass}_R(H^j_I(J^nM/J^{n+1}M))$ with $k = -1$ or $k = 0$, and $\bigcup_{j\leq r_1} \operatorname{Ass}_R(H^j_I(J^nM/J^{n+1}M)) \cup \{\mathfrak{m}\}$ are stable for large n. We also obtain similar results for modules M/J^nM.

1 Introduction

Let $(R, \mathfrak{m})$ be a Noetherian local ring, I, J two ideals of R and M a finitely generated $R-$module. This paper is concerned the asymptotic stability of sets of associated prime ideals $\operatorname{Ass}_R(H^j_I(J^nM/J^{n+1}M))$ and $\operatorname{Ass}_R(H^j_I(M/J^nM))$. There are two starting points for our investigation as follows.

Firstly, In 1979, M. Brodmann [1] had proved that the sets $\operatorname{Ass}_R(J^nM/J^{n+1}M)$ and $\operatorname{Ass}_R(M/J^nM)$ are stable for large n. Based on this result he showed in [2, Theorem 2 and Proposition12] that the integers $\operatorname{depth}(I, J^nM/J^{n+1}M)$ and $\operatorname{depth}(I, M/J^nM)$ take constant values for large n. Recently, M. Brodmann and L.T. Nhan introduced the notion of $M-$sequence in dimension $> k$ in [4]: Let k be an integer $k \geq -1$. A sequence $x_1, \ldots, x_r$ of elements of $\mathfrak{m}$ is called an $M-$sequence in dimension $> k$ if $x_i \notin \mathfrak{p}$ for all $\mathfrak{p} \in \operatorname{Ass}_R(M/(x_1, \ldots, x_{i-1})M)$ with $\dim(R/\mathfrak{p}) > k$ and all $i = 1, \ldots, r$. They also showed that every maximal $M-$sequence in dimension $> k$ in I have the same length,

---

[1] *Key words and phrases*: associated prime, local cohomology, depth, filter depth, generalized depth
2000 Subject Classification: 13D45, 13E05

This work is supported in part by the National Basis Research Programme in Natural Science of Vietnam.

[a] E-mail: ntcuong@math.ac.vn
[b] E-mail: nguyenvanhoang1976@yahoo.com
[c] E-mail: huukhanh75@yahoo.com

and the common length is the least integer i such that there exists $\mathfrak{p} \in \operatorname{Supp}(H_I^i(M))$ with $\dim(R/\mathfrak{p}) > k$. For convenience, we denote this common length by $\operatorname{depth}_k(I, M)$. It should be mentioned that $\operatorname{depth}_{-1}(I, M)$ is just $\operatorname{depth}(I, M)$ the length of a maximal M-sequence in I, $\operatorname{depth}_0(I, M)$ is the filter depth f-depth(I, M) with respect to the ideal I defined by R. Lü and Z. Tang [14], and $\operatorname{depth}_1(I, M)$ is the generalized depth gdepth(I, M) with respect to the ideal I defined by L.T. Nhan [20]. The first result of this paper is a generalization of Brodmann's Theorem.

Theorem 1.1. *For all $k \geq -1$, $\operatorname{depth}_k(I, J^nM/J^{n+1}M)$ and $\operatorname{depth}_k(I, M/J^nM)$ take constant values r_k and s_k, respectively, for large n .*

Secondly, in 1990, C. Huneke [9, Problem 4] asked whether the set of associated primes of $H_I^i(M)$ is finite for all generated modules M and all ideals I. Affirmative answers were given by G. Lyubeznik [15] and Huneke-R.Y. Sharp [10] for equicharacteristic regular local rings. Although, A. Singh [21] and M. Katzman [11] provided examples of finitely generated modules having some local cohomology with infinite associated prime ideals, the problem is still true in many situations (see [3] [12], [13], [16], [20]...). Especially, it was proved in [13] and [20] that $\operatorname{Ass}_R(H_I^j(M))$ is finite for all $j \leq \operatorname{depth}_1(I, M)$. From this and Theorem 1.1 we obtain for all $j \leq r_1 = \operatorname{depth}_1(I, J^nM/J^{n+1}M)$ and $i \leq s_1 = \operatorname{depth}_1(I, M/J^nM)$ that the sets $\operatorname{Ass}_R(H_I^j(J^nM/J^{n+1}M)$ and $\operatorname{Ass}_R(H_I^i(M/J^nM)$ are finite for large n. So it is natural to ask the following question:
Are the sets $\operatorname{Ass}_R(H_I^j(J^nM/J^{n+1}M))$ and $\operatorname{Ass}_R(H_I^i(M/J^nM))$ for $j \leq r_1$ and $i \leq s_1$ stable for large n?

The main result of this paper is the following theorem which shows that a weakening of the above questions may have a positive answer.

Theorem 1.2. *Let $k \geq -1$ be an integer, and r_k, s_k as in Theorem 1.1. Then the following statements are true.*

(i) $\operatorname{Ass}_R(H_I^{r_{-1}}(J^nM/J^{n+1}M))$ *and* $\operatorname{Ass}_R(H_I^{s_{-1}}(M/J^nM))$ *are stable for large n.*

(ii) $\bigcup\limits_{j \leq r_0} \operatorname{Ass}_R(H_I^j(J^nM/J^{n+1}M))$ *and* $\bigcup\limits_{i \leq s_0} \operatorname{Ass}_R(H_I^i(M/J^nM))$ *are stable for large n.*

(iii) $\bigcup\limits_{t \leq j} \operatorname{Ass}_R(H_I^t(J^nM/J^{n+1}M)) \cup \{\mathfrak{m}\}$ *and* $\bigcup\limits_{t \leq i} \operatorname{Ass}_R(H_I^t(M/J^nM)) \cup \{\mathfrak{m}\}$ *for all $j \leq r_1$ and $i \leq s_1$ are stable for large n.*

Notice here that since $r_{-1} \leq r_0 \leq r_1$ as well as $s_{-1} \leq s_0 \leq s_1$ and the modules $H_I^j(J^nM/J^{n+1}M)$, $H_I^i(M/J^nM)$ are either vanished or Artinian for all $j < r_0$ and $i < s_0$ by [14], the question is trivial at these j, i. So the conclusion (i) of Theorem 1.2 is the first non-trivial answer to the above question at $j = r_{-1}$ and $i = s_{-1}$ when $r_{-1} = r_0$ and $s_{-1} = s_0$. Moreover, the conclusion (ii) (the conclusion (iii), respectively) of Theorem 1.2 says that except the maximal ideal $\mathfrak{m}$ (a finite set, respectively) the sets in the question above are stable for all $j \leq r_0$ and $i \leq s_0$ ($j \leq r_1$ and $i \leq s_1$, respectively).

This paper is divided into 5 sections. In Section 2 we study some properties concerning generalizations of the depth of a finite generated module over a local ring. Theorem

1.1 will be also showed in this section. Sections 3, 4 and 5 are devoted to prove the statements (i), (ii) and (iii), respectively, of Theorem 1.2.

2 Asymptotic stability of generalizations of depth

Throughout this paper, $(R, \mathfrak{m})$ is a Noetherian local ring with the maximal ideal $\mathfrak{m}$, I, J two ideals of R and M is a finitely generated $R-$module.

Definition 2.1. ([4, Definition 2.1]) Let $k \geq -1$ be an integer. A sequence $x_1, \ldots, x_r$ of elements of $\mathfrak{m}$ is called an *$M-$sequence in dimension* $> k$ if $x_i \notin \mathfrak{p}$ for all $\mathfrak{p} \in \operatorname{Ass}_R(M/(x_1, \ldots, x_{i-1})M)$ with $\dim(R/\mathfrak{p}) > k$ and all $i = 1, \ldots, r$.

It is clear that $x_1, \ldots, x_r$ is an $M-$sequence in dimension > -1 if and only if it is a regular sequence of M; and $x_1, \ldots, x_r$ is an $M-$sequence in dimension > 0 if and only if it is a filter regular sequence of M introduced by P. Schenzel, N. V. Trung and the first author in [8]. Moreover, $x_1, \ldots, x_r$ is an $M-$sequence in dimension > 1 if and only if it is a generalized regular sequence of M defined by L. T. Nhan in [20].

Remark 2.2. (i) Let k be a non negative integer. Assume that $\dim(M/IM) > k$. Then any $M-$sequence in dimension $> k$ in I is of finite length, and all maximal $M-$sequence in dimension $> k$ in I have the same length which is equal to the least integer i such that there exists $\mathfrak{p} \in \operatorname{Supp}(H_I^i(M))$ with $\dim(R/\mathfrak{p}) > k$ ([4, Lemma 2.4]). We denote in this case by $\operatorname{depth}_k(I, M)$ the length of a maximal $M-$sequence in dimension $> k$ in I. Moreover, if $x_1, \ldots, x_r$ is a maximal $M-$sequence in dimension $> k$ in I, then $x_1, \ldots, x_r$ is a part of system of parameters of M, and thus $\operatorname{depth}_k(I, M) \leq \dim(M) - \dim(M/IM)$. Note that $\operatorname{depth}_{-1}(I, M)$ is the usual depth $\operatorname{depth}(I, M)$ of M in I, $\operatorname{depth}_0(I, M)$ is the filter depth f-depth(I, M) of M in I denoted by Lü and Tang in [14] and $\operatorname{depth}_1(I, M)$ is just the generalized depth gdepth(I, M) of M in I defined by Nhan [20].

(ii) If $\dim(M/IM) \leq k$, then we can choose an $M-$sequence in dimension $> k$ in I of length r for every positive integer r, and we set in this case that $\operatorname{depth}_k(I, M) = \infty$.

Let S be a subset of $\operatorname{Spec}(R)$ and $i \geq 0$ an integer, we set

$$S_{\geq i} = \{\mathfrak{p} \in S | \dim(R/\mathfrak{p}) \geq i\} \text{ and } S_{>i} = \{\mathfrak{p} \in S | \dim(R/\mathfrak{p}) > i\}.$$

Lemma 2.3. *Let $k \geq -1$ be an integer. Then*

$$\begin{aligned}\operatorname{depth}_k(I, M) &= \inf\{j \mid \dim(\operatorname{Ext}_R^j(R/I, M)) > k\} \\ &= \inf\{\operatorname{depth}_{k-i}(I_{\mathfrak{p}}, M_{\mathfrak{p}}) \mid \mathfrak{p} \in \operatorname{Supp}(M/IM)_{\geq i}\}\end{aligned}$$

for all $0 \leq i \leq k+1$, where we use the convenience that $\inf(\emptyset) = \infty$.

Proof. Since $\operatorname{depth}_k(I, M) = \infty$ if and only if $\dim(M/IM) \leq k$, the statement is clear for the case $\operatorname{depth}_k(I, M) = \infty$. Assume that $r = \operatorname{depth}_k(I, M)$ is a non negative integer. By [4, Lemma 2.4], we get

$$r = \inf\{i \mid \exists \mathfrak{p} \in \operatorname{Supp}(H_I^i(M)), \dim(R/\mathfrak{p}) > k\}.$$

Moreover, by [7, Lemma 2.8], we have

$$\bigcup_{j\leq l}\operatorname{Supp}(H_I^j(M))=\bigcup_{j\leq l}\operatorname{Supp}(\operatorname{Ext}_R^j(R/I,M))$$

for all $l\geq 0$. It follows that $r=\inf\{j\mid \dim(\operatorname{Ext}_R^j(R/I,M))>k\}$. To prove the second equality, let $x_1,\ldots,x_r\in I$ be an $M-$sequence in dimension $>k$, and $i\in\{0,\ldots,k+1\}$. For each $\mathfrak{p}\in\operatorname{Supp}(M/IM)_{\geq i}$, $x_1/1,\ldots,x_r/1$ is an $M_{\mathfrak{p}}-$sequence in dimension $>k-i$ in $I_{\mathfrak{p}}$; and thus $r\leq \operatorname{depth}_{k-i}(I_{\mathfrak{p}},M_{\mathfrak{p}})$. On the other hand, by the first equality, there exists $\mathfrak{q}\in\operatorname{Supp}(\operatorname{Ext}_R^r(R/I,M))$ with $\dim(R/\mathfrak{q})>k$. Hence, we can choose a prime ideal $\mathfrak{p}'$ such that $\mathfrak{p}'\supseteq\mathfrak{q}$, $\dim(R/\mathfrak{p}')=i$ and $\dim(R_{\mathfrak{p}'}/\mathfrak{q}R_{\mathfrak{p}'})>k-i$. Therefore $\mathfrak{q}R_{\mathfrak{p}'}\in\operatorname{Supp}_{R_{\mathfrak{p}'}}(\operatorname{Ext}^r_{R_{\mathfrak{p}'}}(R_{\mathfrak{p}'}/I_{\mathfrak{p}'},M_{\mathfrak{p}'}))_{>k-i}$. It follows that $r\geq\operatorname{depth}_{k-i}(I_{\mathfrak{p}'},M_{\mathfrak{p}'})$, and so $r=\inf\{\operatorname{depth}_{k-i}(I_{\mathfrak{p}},M_{\mathfrak{p}})\mid\mathfrak{p}\in\operatorname{Supp}(M/IM)_{\geq i}\}$ as required. ☐

Let $\mathcal{R}=\oplus_{n\geq 0}R_n$ be finitely generated a standard graded algebra over $R_0=R$, and $\mathcal{M}=\oplus_{n\geq 0}M_n$ is a finitely generated graded $\mathcal{R}-$module. For the convenience, we use N_n to denote from now on either the $R-$module M_n or M/J^nM. Then Brodmann's Theorem on the asymptotic stability of associated prime ideals [1] (see also [17, Theorem 3.1]) can be stated as follows.

Lemma 2.4. *The set* $\operatorname{Ass}_R(N_n)$ *is stable for large* n.

Lemma 2.5. *Let* $k\geq -1$ *and* $r\geq 1$ *be integers. If* $\dim(\operatorname{Ext}_R^i(R/I,N_n))\leq k$ *for infinitely many* n *and all* $i<r$. *Then there always exists a sequence of* r *elements* $x_1,\ldots,x_r\in I$ *which is an* N_n-*sequence in dimension* $>k$ *for all large* n.

Proof. Assume that T is an infinite subset of the set of integers such that for all $n\in T$ and $i<r$ we have $\dim(\operatorname{Ext}_R^i(R/I,N_n))\leq k$. We proceed by induction on r. If $r=1$, then $\dim(\operatorname{Hom}(R/I,N_n))\leq k$ for all $n\in T$. It implies $I\nsubseteq\mathfrak{p}$ for all $\mathfrak{p}\in\operatorname{Ass}_R(N_n)_{>k}$ and all $n\in T$. Since $\operatorname{Ass}_R(N_n)$ is stable for large n by Lemma 2.4, there exists an integer $a\in T$ such that $I\nsubseteq\mathfrak{p}$ for all $\mathfrak{p}\in\operatorname{Ass}_R(N_n)_{>k}$ and all $n\geq a$. Hence, we can choose an element $x_1\in I$ which is an N_n-sequence in dimension $>k$ for all $n\geq a$. Assume that $r>1$. Note that $\dim(0:_{N_n}x_1)\leq k$ and $\dim(\operatorname{Ext}_R^i(R/I,N_n))\leq k$ for all $n\in T$ with $n\geq a$ and all $i<r$. From this, by using the following exact sequences

$$\operatorname{Ext}_R^i(R/I,N_n)\to\operatorname{Ext}_R^i(R/I,N_n/(0:_{N_n}x_1))\to\operatorname{Ext}_R^{i+1}(R/I,(0:_{N_n}x_1))$$

and

$$\operatorname{Ext}_R^{i-1}(R/I,N_n)\to\operatorname{Ext}_R^{i-1}(R/I,N_n/x_1N_n)\to\operatorname{Ext}_R^{i}(R/I,N_n/(0:_{N_n}x_1)),$$

we obtain that $\dim(\operatorname{Ext}_R^i(R/I,N_n/x_1N_n))\leq k$ for all $n\in T$ with $n\geq a$ and all $i<r-1$. Hence, by inductive hypothesis, there exists an N_n/x_1N_n-sequence in dimension $>k$ $x_2,\ldots,x_r\in I$ for all $n\geq a$ for some $a\in T$ as required. ☐

Proof of Theorem 1.1. By Lemma 2.4, there is an integer $u>0$ such that $d=\dim(N_n)$ and $d'=\dim(N_n/IN_n)$ for all $n\geq u$. If $d'\leq k$, then $\operatorname{depth}_k(I,N_n)=\infty$ for

$n \geq u$, and the conclusion follows.
Assume that $d' > k$. Then $0 \leq \operatorname{depth}_k(I, N_n) \leq d - d'$ for all $n \geq u$ by Remark 2.2, (i). Thus there exists an infinite subset T of $\mathbb{Z}$, and an integer $r \in \{0, \ldots, d - d'\}$ such that

$$r = \inf\{i \mid \dim(\operatorname{Ext}^i_R(R/I, N_n)) > k\}$$

for all $n \in T$. We will show that $\operatorname{depth}_k(I, N_n) = r$ for all n large. Indeed, if $r = 0$, then $\dim(\operatorname{Hom}(R/I, N_n)) > k$ for all $n \in T$. Since $\operatorname{Ass}_R(\operatorname{Hom}(R/I, N_n)) = \operatorname{Ass}_R(N_n) \cap V(I)$, we get by Lemma 2.4 that the set $\operatorname{Ass}_R(\operatorname{Hom}(R/I, N_n))$ is stable for all $n \geq a$ for some $a \in T$. It follows that $\operatorname{depth}_k(I, N_n) = 0$ for all $n \geq a$. If $r \geq 1$, then we get by Lemma 2.5 that $\operatorname{depth}_k(I, N_n) \geq r$ for all $n \geq v$ for some $v > 0$. Therefore $\dim(\operatorname{Ext}^i_R(R/I, N_n)) \leq k$ for all $n \geq v$ and all $i < r$. Assume that $\dim(\operatorname{Ext}^r_R(R/I, N_n)) \leq k$ for all n in an infinite subset S of $\mathbb{Z}$. Then there exists by Lemma 2.5 an integer $b \geq v$ such that $\operatorname{depth}_k(I, N_n) \geq r + 1$, and so $\dim(\operatorname{Ext}^r_R(R/I, N_n)) \leq k$ for all $n \geq b$ by Lemma 2.3. This contradicts the definition of r. Thus there exists an enough large integer c such that $\dim(\operatorname{Ext}^r_R(R/I, N_n)) > k$ and $\dim(\operatorname{Ext}^i_R(R/I, N_n)) \leq k$ for all $n \geq c$ and $i < r$. Therefore by Lemma 2.3 again we get $\operatorname{depth}_k(I, N_n) = r$ for all $n \geq c$. ☐

In view of Theorem 1.1, if $r = \operatorname{depth}_k(I, N_n)$ for all n large, we call that r is the eventual value of $\operatorname{depth}_k(I, N_n)$. By Lemma 2.5 and the proof of Theorem 1.1, we get immediately the following consequence.

Corollary 2.6. *Let $k \geq -1$ be an integer and r the eventual value of* $\operatorname{depth}_k(I, N_n)$. *Then*

$$r = \inf\{i \mid \dim(\operatorname{Ext}^i_R(R/I, N_n)) > k \text{ for infinitely many } n\}.$$

Lemma 2.7. *Let $k \geq -1$ be an integer. Assume that r and s are the eventual values of* $\operatorname{depth}_k(I, J^nM/J^{n+1}M)$ *and* $\operatorname{depth}_k(I, M/J^nM)$*, respectively. Then we have $r \geq s$.*

Proof. Let $n > 0$. Set $r(n) = \operatorname{depth}_k(I, J^nM/J^{n+1}M)$ and $s(n) = \operatorname{depth}_k(I, M/J^nM)$. By the short exact sequence

$$0 \to J^nM/J^{n+1}M \to M/J^{n+1}M \to M/J^nM \to 0,$$

we get the long exact sequence

$$\cdots \to \operatorname{Ext}^{j-1}_R(R/I, M/J^nM) \to \operatorname{Ext}^j_R(R/I, J^nM/J^{n+1}M) \\ \to \operatorname{Ext}^j_R(R/I, M/J^{n+1}M) \to \cdots .$$

Therefore for any $j < \min\{s(n)+1, s(n+1)\}$ we get by Lemma 2.3 that

$$\dim(\operatorname{Ext}^j_R(R/I, J^nM/J^{n+1}M)) \leq k.$$

It implies that $r(n) \geq \min\{s(n)+1, s(n+1)\}$ by Lemma 2.3. Note that, by the hypothesis of r and s, there exists an integer $a > 0$ such that $r = r(n)$ and $s = s(n)$ for all $n \geq a$. It follows that $r \geq s$. ☐

Recall that a sequence $x_1, \ldots, x_r \in I$ is called an *$I-$filter regular sequence* of M if $x_i \notin \mathfrak{p}$ for all $\mathfrak{p} \in \operatorname{Ass}_R(M/(x_1, \ldots, x_{i-1})M) \setminus V(I)$ and all $i = 1, \ldots, r$, where $V(I)$ is the set of all prime ideals of R containing I.

Lemma 2.8. *Let $k \geq -1$ be an integer and r the eventual value of $\operatorname{depth}_k(I, N_n)$. Assume that $1 \leq r < \infty$. Then there is a sequence $x_1, \ldots, x_r$ in I which is at the same time a permutable N_n-sequence in dimension $> k$ and a permutable $I-$filter regular sequence of N_n for all large n.*

Proof. Let l be an integer that such $1 \leq l \leq r$. We will show by induction on l the existence of a sequence of l elements in I satisfying the conclusion of the lemma. If $l = 1$, then by Lemma 2.4 we may assume that $\operatorname{Ass}_R(N_n)$ is stable for all $n \geq t$ for some integer $t > 0$. Since $r \geq 1$, $I \nsubseteq \mathfrak{p}$ for all $\mathfrak{p} \in \operatorname{Ass}_R(N_n)_{>k}$ and all $n \geq t$. From this, by the Prime Avoidance Theorem, $I \nsubseteq \mathfrak{p}$ for all $\mathfrak{p} \in \operatorname{Ass}_R(N_n)_{>k} \cup (\operatorname{Ass}_R(N_n) \setminus V(I))$ and all $n \geq t$. Therefore, there is an element $x_1 \in I$ satisfying our lemma.

Assume that $1 < l \leq r$ and there already exists a sequence $x_1, \ldots, x_{l-1} \in I$ which is at the same time a permutable N_n-sequence in dimension $> k$ and a permutable $I-$filter regular sequence of N_n for all $n \geq t$ for some integer $t > 0$. Let Ω be the set of all subsets of $\{1, \ldots, l-1\}$. Hence, since Ω is finite, we get by Lemma 2.4 that the set

$$T = \bigcup_{\Lambda \in \ \Omega} \big(\operatorname{Ass}_R(N_n/\Sigma_{j\in\Lambda} x_j N_n)_{>k} \cup (\operatorname{Ass}_R(N_n/\Sigma_{j\in\Lambda} x_j N_n) \setminus V(I))\big)$$

is finite and stable for all n large. Hence there exists an element $x_l \in I$ such that $x_l \notin \mathfrak{p}$ for all $\mathfrak{p} \in T$ by the Prime Avoidance Theorem. Then with the same method that used in the proof of Brodmann-Nhan in [4, Proposition 2.5], we can show that $x_1, \ldots, x_l$ is at the same time a permutable N_n-sequence in dimension $> k$ and a permutable $I-$filter regular sequence of N_n for all large n. ☐

Finally, we recall the following fact which will be used in the sequel.

Lemma 2.9. (see [12, 1.2], [19, 3.4]) *If $x_1, \ldots, x_r$ is an $I-$filter regular sequence of M then we have*

$$H^j_I(M) = \begin{cases} H^j_{(x_1,\ldots,x_r)}(M) & \text{if } j < r \\ H^{j-r}_I(H^r_{(x_1,\ldots,x_r)}(M)) & \text{if } j \geq r. \end{cases}$$

3 Proof of Theorem 1.2, (i)

In the rest of this paper, instead of $\operatorname{depth}_{-1}(I, M)$, $\operatorname{depth}_0(I, M)$ and $\operatorname{depth}_1(I, M)$ we denote, by tradition, by $\operatorname{depth}(I, M)$, f-depth$(I, M)$ and gdepth(I, M), respectively. We first recall a known fact on local cohomology modules.

Lemma 3.1. (cf. [6, Theorem 2.4]) *Let $r = \operatorname{depth}(I, M)$. Assume that $1 \leq r < \infty$ and $x_1, \ldots, x_r$ a regular sequence of M in I. Then we have*

$$\operatorname{Ass}_R(H^r_I(M)) = \operatorname{Ass}_R(M/(x_1, \ldots, x_r)M) \cap V(I).$$

The statement (i) of Theorem 1.2 is then an immediate consequence of following theorems.

Theorem 3.2. *Let r be the eventual value of* $\operatorname{depth}(I, M_n)$*. Then the set* $\operatorname{Ass}_R(H_I^r(M_n))$ *is stable for large n.*

Proof. If $r = \infty$ then $\operatorname{Ass}_R(H_I^r(M_n)) = \emptyset$ for all large n. If $r = 0$ then $\operatorname{Ass}_R(H_I^0(M_n)) = \operatorname{Ass}_R(M_n) \cap V(I)$ is stable for large n by Lemma 2.4. It remains to consider the case $1 \leq r < \infty$. Then we get from Theorem 1.1, Lemma 2.4 and Lemma 2.8 that there exists a positive integer a such that for all $n \geq a$ the following are true: (i) $r = \operatorname{depth}(I, M_n)$; (ii) there exists a sequence $x_1, \ldots, x_r \in I$ which is a regular sequence of M_n; and (iii) the set $\operatorname{Ass}_R(M_n/(x_1, \ldots, x_r)M_n)$ is independent of n. Therefore, we get by Lemma 3.1 that

$$\operatorname{Ass}_R(H_I^r(M_n)) = \operatorname{Ass}_R(M_n/(x_1, \ldots, x_r)M_n) \cap V(I)$$

for all $n \geq a$, and so that $\operatorname{Ass}_R(H_I^r(M_n))$ is stable for large n. ☐

Theorem 3.3. *Let s be the eventual value of* $\operatorname{depth}(I, M/J^nM)$*. Then the set* $\operatorname{Ass}_R(H_I^s(M/J^nM))$ *is stable for large n.*

Proof. Let r be the eventual value of $\operatorname{depth}(I, J^nM/J^{n+1}M)$. Then $r \geq s$ by Lemma 2.7. Moreover, we can choose an integer $a > 0$ such that $r = \operatorname{depth}(I, M/J^nM)$ and $s = \operatorname{depth}(I, J^nM/J^{n+1}M)$ for all $n \geq a$. From the short exact sequence

$$0 \to J^nM/J^{n+1}M \to M/J^{n+1}M \to M/J^nM \to 0,$$

we get the following long exact sequence

$$\cdots \to H_I^{j-1}(M/J^nM) \to H_I^j(J^nM/J^{n+1}M) \to$$
$$\to H_I^j(M/J^{n+1}M) \to H_I^j(M/J^nM) \to \cdots$$

for all $n > 0$. We consider three cases as follows.

Case 1: $r - 2 \geq s$. By the long exact sequence, we have the following isomorphisms

$$H_I^s(M/J^{n+1}M) \cong H_I^s(M/J^nM)$$

for all $n \geq a$. Thus $\operatorname{Ass}_R(H_I^s(M/J^nM))$ is stable for large n.

Case 2: $r - 1 = s$. By the long exact sequence above, we get the exact sequence

$$0 \to H_I^s(M/J^{n+1}M) \to H_I^s(M/J^nM) \to H_I^{s+1}(J^nM/J^{n+1}M)$$

for all $n \geq a$. So $\operatorname{Ass}_R(H_I^s(M/J^{n+1}M)) \subseteq \operatorname{Ass}_R(H_I^s(M/J^nM))$ for all $n \geq a$. It follows that $\operatorname{Ass}_R(H_I^s(M/J^nM))$ is stable for large n, since $\operatorname{Ass}_R(H_I^s(M/J^aM))$ is a finite set by Lemma 3.1.

Case 3: $r = s$. By the long exact sequence, we have the following exact sequence

$$0 \to H_I^s(J^nM/J^{n+1}M) \to H_I^s(M/J^{n+1}M) \to H_I^s(M/J^nM)$$

for all $n \geq a$. There exists by Theorem 3.2 an integer $b \geq a$ such that

$$\mathrm{Ass}_R(H^s_I(J^nM/J^{n+1}M) = \mathrm{Ass}_R(H^s_I(J^bM/J^{b+1}M)$$

for all $n \geq b$. Set $X = \mathrm{Ass}_R(H^s_I(J^bM/J^{b+1}M))$, then we get by the previous exact sequence that

$$X \subseteq \mathrm{Ass}_R(H^s_I(M/J^{n+1}M)) \subseteq \mathrm{Ass}_R(H^s_I(M/J^nM)) \cup X$$

for all $n \geq b$. Hence, for any $n \geq b$, we have

$$\mathrm{Ass}_R(H^s_I(M/J^{n+2}M)) \subseteq \mathrm{Ass}_R(H^s_I(M/J^{n+1}M)) \cup X = \mathrm{Ass}_R(H^s_I(M/J^{n+1}M)).$$

Since $\mathrm{Ass}_R(H^s_I(M/J^{b+1}M))$ is finite by Lemma 3.1, it follows that $\mathrm{Ass}_R(H^s_I(M/J^nM))$ is stable for large n, this finishes the proof of the theorem. □

4 Proof of Theorem 1.2, (ii)

We first need a characterization of filter depth by the Artinianness of local cohomology modules as follows.

Lemma 4.1. ([18, Theorem 3.1]) f-depth$(I, M) = \inf\{i \mid H^i_I(M)$ *is not Artinian* $\}$.

Now, we apply Theorems 3.2 and 3.3 to prove the statement (ii) of Theorem 1.2. We will do this in the two following theorems corresponding to $N_n = M_n$ in the graded case $\mathcal{M} = \oplus_{n\geq 0}M_n$ and $N_n = M/J^nM$.

Theorem 4.2. *Let* r_0 *be the eventual value of* f-depth(I, M_n). *Then the set* $\bigcup\limits_{j\leq r_0} \mathrm{Ass}_R(H^j_I(M_n))$ *is stable for large* n.

Proof. By virtue of Lemmas 2.4 and 4.1 the theorem is trivial for the case of either $r_0 = 0$ or $r = \infty$. Therefore it remains to prove the theorem in the case $1 \leq r_0 < \infty$. Then by Theorem 1.1 and Lemmas 2.4 and 2.8 there exists an integer a such that for all $n \geq a$ the following conditions are true: (i) $r_0 =$ f-depth(I, M_n); (ii) there is a sequence $x_1, \ldots, x_{r_0} \in I$ which is at the same time a permutable filter regular sequence of M_n and a permutable $I-$filter regular sequence of M_n; and (iii) the set $\mathrm{Ass}_R(M_n/(x_1, \ldots, x_{r_0})M_n)$ is independent of n. We first prove that $\bigcup_{n\geq a} \mathrm{Ass}_R(H^{r_0}_I(M_n))$ is a finite set. Let $n \geq a$. We have $H^{r_0}_I(M_n) \cong H^0_I(H^{r_0}_{(x_1,\ldots,x_{r_0})}(M_n)$ by Lemma 2.9, and

$$H^{r_0}_{(x_1,\ldots,x_{r_0})}(M_n) \cong \varinjlim_t (M_n/(x_1^t, \ldots, x_{r_0}^t)M_n)$$

by [5, Theorem 5.2.9]. Thus by virtue of [4, Proposition 2.6] we obtain

$$\mathrm{Ass}_R(H^{r_0}_I(M_n)) \subseteq \bigcup_{t>0} \mathrm{Ass}_R(M_n/(x_1^t, \ldots, x_{r_0}^t)M_n) = \mathrm{Ass}_R(M_n/(x_1, \ldots, x_{r_0})M_n).$$

Therefore the set $\bigcup_{n\geq a} \mathrm{Ass}_R(H^{r_0}_I(M_n))$ is finite by the choice of a. From this we may assume without loss of generality that a large enough such that the set $S =$

$\bigcup_{n\geq a}\operatorname{Ass}_R(H_I^{r_0}(M_n))$ only consists of all prime ideals $\mathfrak{p}$ which belong to $\operatorname{Ass}_R(H_I^{r_0}(M_n))$ for infinitely many $n \geq a$.
Next, we claim that $\operatorname{Ass}_R(H_I^{r_0}(M_n)) \setminus \{\mathfrak{m}\}$ is stable for large n. Indeed, for any $\mathfrak{p} \in S$ there is an infinite set T of integers such that $\mathfrak{p} \in \operatorname{Ass}_R(H_I^{r_0}(M_n))$ for all $n \in T$. Hence $H_{I_\mathfrak{p}}^{r_0}((M_n)_\mathfrak{p}) \neq 0$ so that $\operatorname{depth}(I_\mathfrak{p}, (M_n)_\mathfrak{p}) \leq r_0$. As $\mathfrak{p} \in \operatorname{Supp}(M_n/IM_n) \setminus \{\mathfrak{m}\}$, $\operatorname{depth}(I_\mathfrak{p}, (M_n)_\mathfrak{p}) \geq r_0$ by Lemma 2.3. Thus $\operatorname{depth}(I_\mathfrak{p}, (M_n)_\mathfrak{p}) = r_0$ for all $n \in T$. Keep in mind that $\operatorname{depth}(I_\mathfrak{p}, (M_n)_\mathfrak{p})$ is stable for large n by Theorem 1.1. Hence $\operatorname{depth}(I_\mathfrak{p}, (M_n)_\mathfrak{p}) = r_0$ for all large n. It follows by Theorem 3.2 that $\operatorname{Ass}_{R_\mathfrak{p}}(H_{I_\mathfrak{p}}^{r_0}((M_n)_\mathfrak{p}))$ is stable for large n. Therefore $\operatorname{Ass}_R(H_I^{r_0}(M_n)) \setminus \{\mathfrak{m}\}$ is stable for large n, and the claim follows.
Let r be the eventual value of $\operatorname{depth}(I, M_n)$. It is clear that $r \leq r_0$. If $r = r_0$, then $\bigcup_{j\leq r_0}\operatorname{Ass}_R(H_I^j(M_n)) = \operatorname{Ass}_R(H_I^{r_0}(M_n))$, and this set is stable for large n by Theorem 3.2. If $r < r_0$, there exists an enough large n such that $H_I^r(M_n) \neq 0$, which is an Artinian module by Lemma 4.1. It follows that

$$\bigcup_{j\leq r_0}\operatorname{Ass}_R(H_I^j(M_n)) = \operatorname{Ass}_R(H_I^{r_0}(M_n)) \cup \{\mathfrak{m}\}$$

is stable for large n, and the theorem is proved. ☐

Before proving the statement (ii) of Theorem 1.2 for M/J^nM, we need the following lemma.

Lemma 4.3. *Let A be a submodule of an R-module K. Then we have*

$$\operatorname{Ass}_R(K/A) \setminus \operatorname{Supp}(A) = \operatorname{Ass}_R(K) \setminus \operatorname{Supp}(A).$$

Proof. Let $\mathfrak{p} \in \operatorname{Ass}_R(K/A) \setminus \operatorname{Supp}(A)$ then $\mathfrak{p} = (A : \nu)_R$ for some $\nu \in K$. It yields that $\mathfrak{p}\nu \subseteq A$. So as $A_\mathfrak{p} = 0$ we have $(\mathfrak{p}\nu)_\mathfrak{p} = 0$. From this, since $\mathfrak{p}\nu$ is a finitely generated R-module, it easy to verify that $\mathfrak{p} \subseteq \operatorname{ann}(r\nu)$ for some $r \notin \mathfrak{p}$. For any $x \in \operatorname{ann}(r\nu)$ then $rx \in \operatorname{ann}(\nu) \subseteq (A : \nu)_R = \mathfrak{p}$; so $x \in \mathfrak{p}$, since $r \notin \mathfrak{p}$. Thus $\mathfrak{p} = \operatorname{ann}(r\nu)$, so that $\mathfrak{p} \in \operatorname{Ass}_R(K)$. It follows that $\operatorname{Ass}_R(K/A) \setminus \operatorname{Supp}(A) \subseteq \operatorname{Ass}_R(K) \setminus \operatorname{Supp}(A)$. The converse inclusion is trivial. ☐

Theorem 4.4. *Let s_0 be the eventual value of* $\text{f-depth}(I, M/J^nM)$. *Then the set* $\bigcup_{j\leq s_0}\operatorname{Ass}_R(H_I^j(M/J^nM))$ *is stable for large n.*

Proof. By Theorem 1.1, there is an integer a such that $r_0 = \text{f-depth}(I, J^nM/J^{n+1}M)$ and $s_0 = \text{f-depth}(I, M/J^nM)$ for all $n \geq a$. By Theorem 3.3 the result is trivial for the case $s_0 = \infty$. Assume now that $s_0 < \infty$. Using the last part in the proof of Theorem 4.2 we need only to show that the set $\operatorname{Ass}_R(H_I^{s_0}(M/J^nM)) \setminus \{\mathfrak{m}\}$ is stable for large n. From the short exact sequence $0 \to J^nM/J^{n+1}M \to M/J^{n+1}M \to M/J^nM \to 0$, we get the following long exact sequence of local cohomology modules

$$\begin{aligned} \cdots \to H_I^{j-1}(M/J^nM) \to H_I^j(J^nM/J^{n+1}M) \\ \to H_I^j(M/J^{n+1}M) \to H_I^j(M/J^nM) \to \cdots \end{aligned}$$

for all $n > 0$. Note that $r_0 \geq s_0$ by Lemma 2.7. Thus we consider two cases as follows.

Case 1: $r_0 - 1 \geq s_0$. By the long exact sequence, we get the following exact sequences

$$H_I^{s_0}(J^nM/J^{n+1}M) \to H_I^{s_0}(M/J^{n+1}M) \to H_I^{s_0}(M/J^nM),$$

for all $n \geq a$, in which $H_I^{s_0}(J^nM/J^{n+1}M)$ is Artinian by Lemma 4.1. Thus we have

$$\mathrm{Ass}_R(H_I^{s_0}(M/J^{n+1}M)) \setminus \{\mathfrak{m}\} \subseteq \mathrm{Ass}_R(H_I^{s_0}(M/J^nM)) \setminus \{\mathfrak{m}\}$$

for all $n \geq a$. Since $\mathrm{Ass}_R(H_I^{s_0}(M/J^nM))$ is finite for $n = a$ by [6, Theorem 2.5], the set $\mathrm{Ass}_R(H_I^{s_0}(M/J^nM)) \setminus \{\mathfrak{m}\}$ is stable for large n.

Case 2: $r_0 = s_0$. Note by Theorem 4.2 that $\mathrm{Ass}_R(H_I^{s_0}(J^nM/J^{n+1}M)) \setminus \{\mathfrak{m}\}$ is stable for all $n \geq b$ for some integer $b \geq a$. Set $X = \mathrm{Ass}_R(H_I^{s_0}(J^bM/J^{b+1}M)) \setminus \{\mathfrak{m}\}$. Thus we get by the exact sequence

$$H_I^{s_0-1}(M/J^nM) \to H_I^{s_0}(J^nM/J^{n+1}M) \to H_I^{s_0}(M/J^{n+1}M) \to H_I^{s_0}(M/J^nM)$$

for all $n \geq a$, in which $H_I^{s_0-1}(M/J^nM)$ is Artinian by Lemma 4.1, and Lemma 4.3 that

$$X \subseteq \mathrm{Ass}_R(H_I^{s_0}(M/J^{n+1}M)) \setminus \{\mathfrak{m}\} \subseteq (\mathrm{Ass}_R(H_I^{s_0}(M/J^nM)) \setminus \{\mathfrak{m}\}) \cup X$$

for all $n \geq b$. Therefore

$$\begin{aligned} \mathrm{Ass}_R(H_I^{s_0}(M/J^{n+2}M)) \setminus \{\mathfrak{m}\} &\subseteq \big(\mathrm{Ass}_R(H_I^{s_0}(M/J^{n+1}M)) \setminus \{\mathfrak{m}\}\big) \cup X \\ &= \mathrm{Ass}_R(H_I^{s_0}(M/J^{n+1}M)) \setminus \{\mathfrak{m}\} \end{aligned}$$

for all $n \geq b$. Note that $\mathrm{Ass}_R(H_I^{s_0}(M/J^{n+1}M)) \setminus \{\mathfrak{m}\}$ is a finite set for $n = b$ by [6, Theorem 2.5]. It follows that $\mathrm{Ass}_R(H_I^{s_0}(M/J^nM)) \setminus \{\mathfrak{m}\}$ is stable for large n as required. ☐

5 Proof of Theorem 1.2, (iii)

To prove the statement (iii) of Theorem 1.2 we first need the following known results.

Lemma 5.1. ([7, Corollary 4.4]) *The following equality holds true*

$$\mathrm{gdepth}(I, M) = \inf\{i \mid \mathrm{Supp}(H_I^i(M)) \textit{ is not finite }\}.$$

Lemma 5.2. ([7, Corollary 4.6]) *The set* $\mathrm{Ass}(H_I^j(N))$ *is finite for all* $j \leq \mathrm{gdepth}(I, N)$.

The first conclution of Theorem 1.2, (iii) is a special case of the following result.

Theorem 5.3. *Let* r_1 *be the eventual value of* $\mathrm{gdepth}(I, M_n)$. *Then for each* $l \leq r_1$ *the set* $\bigcup_{j \leq l} \mathrm{Ass}_R(H_I^j(M_n)) \cup \{\mathfrak{m}\}$ *is stable for large* n.

Proof. By theorem 1.1 and Lemmas 2.4 and 2.8 we can choose a positive integer a such that $d = \dim(M_n)$, $d' = \dim(M_n/IM_n)$ and $r_1 = \operatorname{gdepth}(I, M_n)$ for all $n \geq a$. We first claim that there is an integer $b \geq a$ such that $S = \bigcup_{n \geq b} \bigcup_{j \leq l} \operatorname{Ass}_R(H_I^j(M_n))$ is finite. In order to do this we consider three cases: $r_1 = 0$, $r_1 = \infty$ and $1 \leq r_1 < \infty$. If $r_1 = 0$, then $S \subseteq \bigcup_{n \geq a} \operatorname{Ass}_R(M_n)$ is finite by Lemma 2.4. If $r_1 = \infty$, then $d' \leq 1$, so that $S \subseteq \bigcup_{n \geq a} \operatorname{Supp}_R(M_n/IM_n)$ is finite by Lemma 2.4. Now we consider the case $1 \leq r_1 < \infty$. Then, by Lemma 2.8 we can choose an integer $b \geq a$ such that the following statements are true for all $n \geq b$: (i) there is a sequence $x_1, \ldots, x_{r_1} \in I$ which is at the same time a permutable generalized regular sequence of M_n and a permutable $I-$filter regular sequence of M_n; and (ii) the sets $\operatorname{Ass}_R(M_n/(y_1, \ldots, y_j)M_n)$ are stable for all $j \leq r_1$. From this and [4, Proposition 2.6] we have $S \subseteq \bigcup_{n \geq b} \bigcup_{j \leq l} \operatorname{Ass}_R(M_n/(y_1, \ldots, y_j)M_n)$. Therefore S is finite by Lemma 2.4 again, and the claim is proved.

Since S is finite, we can choose the integer b enough large such that S only consists of all prime ideals $\mathfrak{p}$, which belong to $\bigcup_{j \leq l} \operatorname{Ass}_R(H_I^j(M_n))$ for infinitely many $n \geq b$. For any $\mathfrak{p} \in S_{\geq 1}$, by Theorem 1.1, there is an integer $n(\mathfrak{p}) \geq b$ such that $r(\mathfrak{p}) = \operatorname{depth}(I_{\mathfrak{p}}, (M_n)_{\mathfrak{p}})$ and $r_0(\mathfrak{p}) = \text{f-depth}(I_{\mathfrak{p}}, (M_n)_{\mathfrak{p}})$ for all $n \geq n(\mathfrak{p})$. Then, by the choice of S and Lemma 2.3, we obtain that $r(\mathfrak{p}) \leq l \leq r_1 \leq r_0(\mathfrak{p})$. If $l < r_0(\mathfrak{p})$, $\bigcup_{j \leq l} \operatorname{Ass}_{R_{\mathfrak{p}}}(H_{I_{\mathfrak{p}}}^j((M_n)_{\mathfrak{p}})) = \{\mathfrak{p}R_{\mathfrak{p}}\}$ for all $n \geq n(\mathfrak{p})$ by Lemma 4.1. If $l = r_0(\mathfrak{p})$, it follows from Theorem 4.2 that $\bigcup_{j \leq l} \operatorname{Ass}_{R_{\mathfrak{p}}}(H_{I_{\mathfrak{p}}}^j((M_n)_{\mathfrak{p}}))$ is stable for all $n \geq m(\mathfrak{p})$ for some integer $m(\mathfrak{p}) \geq b$. Therefore $\bigcup_{j \leq l} \operatorname{Ass}_R(H_I^j(M_n)) \cup \{\mathfrak{m}\}$ is stable for all $n \geq \max\{n(\mathfrak{p}),\ m(\mathfrak{p}) \mid \mathfrak{p} \in S_{\geq 1}\}$ as required. ☐

Finally, by applying Theorem 5.3 we can prove the statement (iii) of Theorem 1.2 for the $R-$modules M/J^nM.

Theorem 5.4. *Let s_1 be the eventual value of* $\operatorname{gdepth}(I, M/J^nM)$. *Then for each $l \leq s_1$ the set $\bigcup_{j \leq l} \operatorname{Ass}_R(H_I^j(M/J^nM)) \cup \{\mathfrak{m}\}$ is stable for large n.*

Proof. By Theorem 1.1, there is an integer a such that $r_1 = \operatorname{gdepth}(I, J^nM/J^{n+1}M)$ and $s_1 = \operatorname{gdepth}(I, M/J^nM)$ for all $n \geq a$. Note that $s_1 \leq r_1$ by Lemma 2.7. For any $0 \leq l \leq s_1$ we set

$$X_l(n) = \bigcup_{j \leq l} \operatorname{Ass}_R(H_I^j(M/J^nM)) \cup \{\mathfrak{m}\} \text{ and } S_l(n) = \bigcup_{j \leq l} \operatorname{Ass}_R(H_I^j(J^nM/J^{n+1}M)) \cup \{\mathfrak{m}\}.$$

We will show by induction on l that the set $X_l(n)$ is stable for large n. It is nothing to prove for the case $l = 0$. Let $l > 0$. From the short exact sequence

$$0 \to J^nM/J^{n+1}M \to M/J^{n+1}M \to M/J^nM \to 0$$

we get the following long exact sequence

$$\cdots \to H_I^{j-1}(M/J^nM) \to H_I^j(J^nM/J^{n+1}M) \xrightarrow{f_j} H_I^j(M/J^{n+1}M) \to H_I^j(M/J^nM) \to \cdots$$

for all $n > 0$. From this, for any $j \leq l$ and $n \geq a$ we have

$$\operatorname{Ass}_R(H_I^j(M/J^{n+1}M)) \subseteq \operatorname{Ass}_R(H_I^j(M/J^nM)) \cup \operatorname{Ass}_R(\operatorname{Im}(f_j)).$$

Note that

$$\mathrm{Ass}_R(\mathrm{Im}(f_j)) \subseteq \mathrm{Ass}_R(H_I^j(J^nM/J^{n+1}M)) \cup \mathrm{Supp}_R(\mathrm{Ker}(f_j))$$

by Lemma 4.3. Since $j-1 < s_1$, $\mathrm{Supp}_R(H_I^{j-1}(M/J^nM))$ is a finite set by Lemma 5.1, so that

$$\mathrm{Supp}_R(\mathrm{Ker}(f_j)) \subseteq \mathrm{Supp}_R(H_I^{j-1}(M/J^nM)) \subseteq \mathrm{Ass}_R(H_I^{j-1}(M/J^nM)) \cup \{\mathfrak{m}\}.$$

Therefore we get

$$X_l(n+1) \subseteq X_l(n) \cup S_l(n).$$

On the other hand, by the long exact sequence above and Lemma 5.1, we have

$$\begin{aligned}\mathrm{Ass}_R(H_I^j(J^nM/J^{n+1}M)) &\subseteq \mathrm{Ass}_R(H_I^j(M/J^{n+1}M)) \cup \mathrm{Supp}_R(H_I^{j-1}(M/J^nM)) \\ &\subseteq \mathrm{Ass}_R(H_I^j(M/J^{n+1}M)) \cup \mathrm{Ass}_R(H_I^{j-1}(M/J^nM)) \cup \{\mathfrak{m}\}.\end{aligned}$$

Hence

$$S_l(n) \subseteq X_l(n+1) \cup X_{l-1}(n)$$

for all $n \geq a$. By Theorem 5.3 and the inductive hypothesis, there exists an integer $b \geq a$ such that $S_l(n) = S$ and $X_{l-1}(n) = X$ for all $n \geq b$. Therefore, since $X = X_{l-1}(n+1) \subseteq X_l(n+1)$,

$$S \subseteq X_l(n+1) \cup X = X_l(n+1)$$

for all $n \geq b$. Finally, we obtain the following inclusions

$$X_l(n+2) \subseteq X_l(n+1) \cup S \subseteq X_l(n+1)$$

for all $n \geq b$. Thus $X(n)$ is stable for large n, since $X_l(b+1)$ is finite by Lemma 5.2, and the proof of Theorem 5.4 is complete. ☐

References

[1] M. Brodmann, *Asymptotic stability of* $\mathrm{Ass}_R(M/I^nM)$, Proc. Amer. Math. Soc., (1) **74** (1979), 16 - 18.

[2] M. Brodmann, *The asymptotic nature of the analytic spread*, Math. Proc. Camb. Phil. Soc., **86** (1979), 35 - 39.

[3] M. Brodmann and A. L. Faghani, *A finiteness result for associated primes of local cohomology modules*, Proc. Amer. Math. Soc., (10) **128** (2000), 2851 - 2853.

[4] M. Brodmann and L. T. Nhan, *A finiteness result for associated primes of certain Ext-modules*, to appear in Comm. Algebra.

[5] M. Brodmann and R. Y. Sharp, "Local cohomology: an algebraic introduction with geometric applications", Cambridge University Press, 1998.

[6] N. T. Cuong and N. V. Hoang, *Some finite properties of generalized local cohomology modules,* East-West J. Math., (2) **7** (2005), 107 - 115.

[7] N. T. Cuong and N.V. Hoang, *On the vanishing and the finiteness of supports of generalized local cohomology modules,* to appear in Manuscripta Math., 2008.

[8] N. T. Cuong, P. Schenzel and N. V. Trung, *Verallgemeinerte Cohen-Macaulay Moduln,* Math. Nachr., **85** (1978), 57 - 73.

[9] C. Huneke, *Problems on local cohomology*, Free resolutions in commutative algebra and algebraic geometry (Sundance, Utah, 1990), Res. Notes Math., **2** (1992), 93 - 108.

[10] C. Huneke and R. Y. Sharp, *Bass numbers of local cohomology modules,* Trans. Amer. Math. Soc., **339** (1993), 765 - 779.

[11] M. Katzman, *An example of an infinite set of associated primes of a local cohomology module,* J. Algebra, **252** (2002), 161 - 166.

[12] K. Khashyarmanesh and Sh. Salarian, *Filter regular sequences and the finiteness of local cohomology modules*, Comm. Algebra, (8) **26** (1998), 2483 - 2490.

[13] K. Khashyarmanesh and Sh. Salarian, *On the associated primes of local cohomology modules*, Comm. Algebra, **27** (1999), 6191 - 6198.

[14] R. Lü and Z. Tang, *The f-depth of an ideal on a module,* Proc. Amer. Math. Soc., (7) **130** (2001), 1905 - 1912.

[15] G. Luybeznik, *Finiteness properties of local cohomoly modules (an application of D-modules to commutative algebra),* Invent. Math., **113** (1993), 41 - 55.

[16] T. Marley, *Associated primes of local cohomology module over rings of small dimension,* Manuscripta Math., (4) **104** (2001), 519 - 525.

[17] L. Melkersson, *On asymptotic stability for sets of prime ideals connected with the powers of an ideal*, Math. Proc. Camb. Phil. Soc., **107** (1990), 267 - 271.

[18] L. Melkersson, *Some applications of a criterion for artinianness of module,* J. Pure Appl. Algebra, **101** (1995), 291 - 303.

[19] U. Nagel and P. Schenzel, *Cohomological annihilators and Castelnuovo-Mumford regularity,* In: Commutative Algebra: Syzygies, Multiplicities, and Birational Algebra (South Hadley, 1992). Contemp. Math., 159. Amer. Math. Soc., (1994), 307 - 328.

[20] L. T. Nhan, *On generalized regular sequences and the finiteness for associated primes of local cohomology modules,* Comm. Algebra, **33** (2005), 793 - 806.

[21] A. Singh, *$p-$torsion elements in local cohomology modules*, Math. Res. Lett., **7** (2000), 165 - 176.